

Shufly SM

National Aeronautics and Space Administration
Goddard Space Flight Center
Contract No. NAS-5-9299

ST - CM — 10 462

CONDITION REQUIRED FOR THE STABILITY OF
ORBITAL MOTION

by
K. V. Kholoshevnikov
(USSR)

FACILITY FORM 602

N66-87192
(ACCESSION NUMBER)

4
(PAGES)

CA 78098
(NASA CR OR TMX OR AD NUMBER)

(THRU)

None
(CODE)

(CATEGORY)

23 MARCH 1966

CONDITION REQUIRED FOR THE STABILITY OF ORBITAL MOTION

Vestnik Leningradskogo Universiteta
 Seriya Matematiki, Mekhaniki i Astronomii
 No. 1, vypusk 1, 172-3,
 LENINGRAD, 1966.

by K. V. Kholoshevnikov

SUMMARY

It is proved that to achieve the stability of the orbital motion the condition $h < 0$ is necessary, h being the energy constant.

*
* *

1.- We shall consider the motion of a particle in the gravitational field of an arbitrary body (further called the planet), having an axis-symmetrical structure. By strength of the condition of field conservation there exists the energy integral

$$T = V + h, \quad (1)$$

and the validity of the Laplace formula

$$\frac{d^2 R}{dt^2} = U + 4h. \quad (2)$$

Here h is the energy constant; T is the kinetic energy of the mass unit of the particle; V is a mass-flow function; $U = 4V + 2QV$; $R = r^2$, where r is the radius-vector;

$$Q = x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} + z \frac{\partial}{\partial z}.$$

Utilizing the homogeneity of the mass-flow function of the problem of n bodies, Jacobi derived from the formulas (1) and (2) the necessary condition $h < 0$ of solar system's stability.

* NEOBKHODIMOYE USLOVIYE USTOYCHIVOSTI ORBITAL'NOGO DVIZHENIYA

In our case the mass-flow function

$$V = \kappa^2 \left\{ \frac{1}{r} - \sum_{n=2}^{\infty} J_n \frac{P_n\left(\frac{z}{r}\right)}{r^{n+1}} \right\} \quad (3)$$

is devoid of homogeneity. Here κ^2 is the product of the gravitational constant by the mass of the planet; P_n are the Legendre polynomials; J_n are constants. We took for the unit of length the greatest radius-vector of planet's surface.

According to Lagrange, the motion of a material point is said to be steady, if at $t \geq t_0$ the point does emerge from the finite region of space M .

If we take for the region M a sphere of arbitrary radius, it is well known that the condition

$$h < 0. \quad (4)$$

is sufficient for the stability.

We shall demonstrate that the condition (4) is indispensable if we take for M any closed finite region of space D , which would not intersect the planet, and in which

$$U > 0. \quad (5)$$

Assume $h \geq 0$, and that the particle shall never emerge from the region D . Then, it follows from formulas (2) and (5):

$$\frac{d^2 R}{dt^2} > U_0 > 0. \quad (6)$$

Integrating, we find

$$R - R_0 > \frac{U_0}{2} (t - t_0)^2 + \frac{dR(t_0)}{dt} (t - t_0), \quad (7)$$

whence $R = r^2 \frac{1}{1-\epsilon} \rightarrow \infty$, which is in contradiction with the assumption just made. Thus the requirement of the condition (4) has been demonstrated.

2. - It is quite probable that for planets of the solar system the inequality (5) is fulfilled in the whole outer space. If this is so, then at $h \gg 0$ the particle either drifts away to infinity or falls on the planet.*

* This seems evident. However, it is possible to show that there exist force fields in which the motion along the circle $r = r_0 > 1$ is possible for any great h .

It has been possible to demonstrate mathematically the inequality (5) for $r \gg 1$, for the series (3) may be divergent at $r < 1$. To demonstrate this, let us note first of all that at $r \gg 1$ the series (3) may be differentiated over x, y, z termwise, for the coefficients J_n of the arbitrary body of revolution [1] satisfy the inequality

$$|J_n| \leq \frac{C}{n^{5/2}}, \quad (8)$$

where C is a certain constant. That is why the operator Δ may be introduced under the sign of the sum (3) and we may take advantage of the Euler formula for an homogenous function. As a result we shall obtain

$$U = \frac{2\pi^2}{r} \left\{ 1 + \sum_{n=2}^{\infty} (n-1) J_n \frac{P_n\left(\frac{z}{r}\right)}{r^n} \right\}. \quad (9)$$

Assume $J_k = \sup_{n \geq k} \{|J_n| n^{5/2}\}$ ($J_k \leq C$).

Since

$$\left| \sum_{n=2}^{\infty} (n-1) J_n \frac{P_n}{r^n} \right| < \frac{|J_2|}{r^3} + j_3 \sum_{n=3}^{\infty} \frac{1}{r^n n^{3/2}} < \frac{1}{r^3} \left\{ |J_2| + J_3 \left[\zeta\left(\frac{3}{2}\right) - 1 - \frac{1}{2^{3/2}} \right] \right\} \quad (r \gg 1),$$

the quantity U will be strictly positive at $r \gg 1$ if

$$|J_2| + j_3 \left[\zeta\left(\frac{3}{2}\right) - 1 - \frac{1}{2^{3/2}} \right] = |J_2| + 1.26 j_3 > 1. \quad (10)$$

Here $\zeta(x)$ is a Riemann-zeta-function.

For the Earth [2] $j_3 < J_2 \approx 10^{-3}$. Obviously, the inequality (10) is fulfilled for all the planets of the solar system.

Therefore, the motion of a particle in the gravitational field of an axi-symmetrical planet may, according to Lagrange, take place in the M-region only in the case when the inequality $h < 0$ is satisfied.

We may take for the region M any finite part of space $r \gg 1$, which coincides with any finite part relative to the planet of the outer space, with a precision to the difference between the equatorial and polar radii of the planet.

*** THE END ***

REFERENCES

- [1]. K. V. KHOLSHEVNIKOV.- O velichine koeffitsientov razlozheniya potentsiala Vestn.LGU, 13, 155, 1965
- [2].- Y. KOZAI.- The potential of the earth derived from satellite motions. Dynamics of satellites.- Berlin, 1962 (Springer).

Contract No.NAS-5-9299
Consultants & Designers, Inc.
Arlington, Virginia

Translated by ANDRE L. BRICHANT
on 23 March 1966

DISTRIBUTIONGODDARD SPACE F.C.NASA HQSOTHER CENTERS

100 CLARK, TOWNSEND
110 STROUD
400 BOURDEAU
610 MEREDITH
611 McDONALD
ABRAHAM, BOLDT
VKB, WILLIAMS
SERLEMITSOS
612 HEPPNER
NESS
613 KUPPERIAN
HALLAM
614 WHITE
615 BAUER
GOLDBERG
STONE
640 HESS
MEAD
NORTHROP
CAMERON
MUSEN
643 SQUIRES
630 GI for SS [3]
500 MENGEL
547 SIRY
730 STAMPFL
252 LIBRARY
256 FREAS

SS NEWELL, NAUGLE
SG MITCHELL
SCHARDT
DUBIN
SL LIDDEL
BRYSON
FELLOWS
HIPSHER
HOROWITZ
SM FOSTER
GILL
RR KURZWEG
RRA WILSON
REC GOULD
RTR NEILL
ATSS SCHWIND
ROBBINS
WX SWEET

Ames RC
SONETT
LIBRARY
Langley RC
160 ADAMSON
213 KATZOFF
BUGLIA
304 BIRD
QUEIJO
185 WEATHERWAX
UCLA
KAULA
JPL
WYCKOFF
WPAFB
Code TDBX-T